




Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: [www.elsevier.com/locate/apjtm](http://www.elsevier.com/locate/apjtm)

Document heading

doi: 10.1016/S1995-7645(14)60259-5

# Traditional effects of medicinal plants in the treatment of respiratory diseases and disorders: an ethnobotanical study in the Urmia

Mohsen Asadbeigi<sup>1</sup>, Tahereh Mohammadi<sup>2</sup>, Mahmoud Rafeian-Kopaei<sup>3</sup>, Kourosh Saki<sup>4</sup>, Mahmoud Bahmani<sup>1\*</sup>, Mohammad Delfan<sup>2</sup>

<sup>1</sup>Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Khorramabad, Iran

<sup>2</sup>Tarbiat Modares University, Tehran, Iran

<sup>3</sup>Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

<sup>4</sup>Shahid Beheshti University of Medical Sciences, Tehran, Iran

## ARTICLE INFO

### Article history:

Received 12 Jul 2014

Received in revised form 14 Aug 2014

Accepted 25 Aug 2014

Available online 17 Sep 2014

### Keywords:

Medicinal plants

Respiratory diseases

Urmia

Herbalists

## ABSTRACT

**Objective:** To identify, present and review the respiratory medicinal plants which used by Urmian herbalists.

**Methods:** The list of traditional healers of West Azarbaijan Province was prepared and data were obtained by direct observation, interviews and the questionnaires. After that, herbarium samples were collected from the desired area and deposited in herbarium unit of the Faculty of Agriculture, Urmia University, Urmia, Iran.

**Results:** Our results demonstrated that 20 medicinal plants from 10 plant families are used to treat respiratory disorders. Also, the most plant part that used for treating of respiratory disorders was seed (27%) and the most traditional form prescribed by herbalists was boiled (54%). Forty three percentage of Urmia herbalists have used herbs for the treatment of cough.

**Conclusions:** People in this area have a strong belief that plants have a positive impact in the treatment of respiratory disorders and they have used medicinal plants since ancient times to treat these disorders. Our study revealed the importance of herbal medicines and traditional medicine in this area as medicinal resource for drug discovery in future.

## 1. Introduction

Upper and lower respiratory tract anatomically are divided into two parts which are separated from each other by the throat. Respiratory diseases can involve the respiratory tract, lungs or blood vessels and usually a combination of these abnormalities can be seen in many respiratory diseases[1,2]. Respiratory diseases can generally classified into the following groups[2]: obstructive pulmonary disease, restrictive lung disease, pulmonary vascular disease and other diseases.

Traditional medicine has a long history of serving peoples all over the world. In many countries and cultures of different nations, the use of medicinal plants to treat diseases and maintain public health is highly prevalent[3–6]. Natural products play an important role in the field of new drugs research and development. Recent studies have also revealed promising results from using of plants in the treatment or prevention of a wide variety of hard curable diseases such, atherosclerosis[7,8], diabetes[9,10], cardiovascular diseases[11,12], neurological disorders[13,14], and cancer[15,16]. The main proposed mechanism for beneficial effects of traditional plant is alterations in redox state[17,18]. Therefore, medicinal plants with antioxidant activity have been shown to counteract these situations and always been considered as a healthy source of health promotion. Extraction of medicinal plants, drug preparation,

\*Corresponding author: Dr. Mahmoud Bahmani, Razi Herbal Medicines Research Center, Lorestan University of Medical Sciences, Po.Box: 6813833946, Khorramabad, Iran.

Tel: +98 661 2202033

E-mail: [mahmood.bahmani@gmail.com](mailto:mahmood.bahmani@gmail.com)

Foundation Project: Supported by research grant (No. 453/03) from Deputy for Research and Technology of Lorestan University of Medical Sciences.

local efforts to document valuable informations on folk medicine, effectiveness and safety in respiratory disease treatment are important for the indigenous peoples of West Azerbaijan Province. Therefore, the aims of this study were to gather local knowledge and traditional medicine of ethnobotany of Uremia region to identify herbs that are effective in the treatment of respiratory diseases.


## 2. Materials and methods

In this study the data were collected by interview and questionnaire during the period of April to June 2013 using public resources. List of traditional healers in Urmia City was prepared from Urmia Drug and Food Administration and then necessary information of traditional healers was collected by direct observation, interview and questionnaire methods. Questionnaires were distributed between Urmia traditional healers and at the same time interview was performed. Herbal samples were collected from the desired region and after drying, their herbarium specimens were prepared. The herbarium samples obtained from data of local traditional physicians in the questionnaire were collected from the region and then they were authenticated by a botanist using a variety of flora and valid references. A herbarium specimen from each plant (whole or the used parts) was prepared and deposited in the herbarium unit of Shahrekord University of Medical Sciences, Shahrkord, Iran.

## 3. Results


After analyzing the data obtained from questionnaires and interviews in Uremia, it has been demonstrated that overall, 20 medicinal plants from 10 plant families were used for the treatment of respiratory disease/disorders. The number and composition of plant families used in the treatment of respiratory diseases and disorders in Uremia were presented in Figure 1. These data demonstrated that Asteraceae and Malvaceae composed the most parts of plants which used in folk medicine with 20% and 15%, respectively. Parts of the plants which used in treatment of respiratory disease were presented in Figure 2. These results showed that the most part of plant that used in treatment of respiratory diseases is seed (27%). Cough was the most sign of respiratory diseases that treated by folk medicinal plant (43%). After that, treating of Asthma and stimulating of phlegm were the most therapeutic targets in using of medicinal plants (Figure 3). The forms of plant using in treatment of respiratory diseases were presented in Figure 4 and demonstrated that the most prescribed herbal decoction form, was run by groceries

(54%). Table 1 showed the most characteristics of our plants with their therapeutic properties.




**Figure 1.** The number and composition of plant families used in the treatment of respiratory diseases and disorders in Uremia.

A: Apiaceae; B: Asteraceae; C: Brassicaceae; D: Fabaceae; E: Lamiaceae; F: Malvaceae; G: Papaveraceae; H: Plantaginaceae; I: Platanaceae; J: Rosaceae.




**Figure 2.** Parts of the plants which used in treatment of respiratory disease. FB: Flower branches.



**Figure 3.** The most sign of respiratory diseases that treated by folk medicinal plant.

A: Cough; B: Pulmonary pain; C: Stimulating expectorant; D: Sound jamming; E: Asthma; F: Pulmonary infection; G: Sorethroat.



**Figure 4.** The forms of plant using in treatment of respiratory diseases.

## 4. Discussion

In this study, the ethnobotanic data about medicinal plants which used in folk medicine from different regions

**Table 1**

The characteristics of medicinal plants with their therapeutic properties.

Scientific name	Family	Persian name	Used parts	Type use	Traditional therapeutic effect
<i>Althea hirsuta</i> L.	Malvaceae	Khatmi	Root	Boiled, brewed, incense	Pulmonary infection Inducing mucus Cough
<i>Alyssum desertorum</i> Stapf.	Brassicaceae	Ghodoumeh	Seed	Boiled, brewed, incense	Cough
<i>Astragalus effuses</i>	Fabaceae	Gavan	Resin	Boiled, brewed	Cough
<i>Astragalus gossypinus</i> Fisch.	Fabaceae	Gavan panbei	Resin	Boiled, brewed, incense	Cough
<i>Cardaria draba</i> (L.) Desv.	Brassicaceae	Azmak	Leaf and Seed	Boiled, brewed, incense	Pulmonary infection Inducing mucus Cough
<i>Cotoneaster nummularia</i> F.M.	Rosaceae	Shirkhesht	Areal parts	Boiled	Inducing cough and mucus
<i>Cuminum cyminum</i> L.	Apiaceae	Zire sabz	Seed	Boiled	Asthma
<i>Echinops ritrodes</i> L.	Asteraceae	Shekar tighal	Stem	Boiled	Chronic cough Sore throat
<i>Hibiscus trionum</i> L.	Malvaceae	Khatmi serang	Flower	Boiled, brewed	Cough Inducing mucus
<i>Hypericum perforatum</i> L.	Asteraceae	Alafe chay	Flowering shoot	Boiled	Cough
<i>Malva neglecta</i> Wallr.	Malvaceae	Panirak	Seed, leaf, flowering shoot	Decoction, poultice	Coughing Expectorant
<i>Mentha longifolia</i> L.	Lamiaceae	Pouneh	Areal parts	Boiled, brewed, incense	Pulmonary infection Sore throat Mucus binding Treatment of asthma
<i>Papaver rhoeas</i> L.	Papaveraceae	Shaghayegh	Seed, capsule	Boiled	Inducing mucus Cough
<i>Pimpinella affinis</i> Ledeb.	Apiaceae	Tartizak baghi	Flowering shoot, seed	Boiled	Cough Chest pain relief
<i>Plantago major</i> L.	Plantaginaceae	Barhang	Seed, leaf, root	Boiled	Cough Pulmonary infection
<i>Platanus orientalis</i> L.	Platanaceae	Chenar	Plant skin	Boiled	Treatment of asthma Hoarseness
<i>Rubus persicum</i> Boiss.	Rosaceae	Tameshk	Fruit	Boiled, raw edible	Asthma
<i>Scrozonera cinerea</i> Boiss.	Asteraceae	Gounei sheng	Root	Boiled	Cough
<i>Thymus kotschyanus</i> Boiss.	Lamiaceae	Avishan	Flowering shoot	Brewed, incense	Cough, Asthma
<i>Tragopogon carcifolius</i> Boiss.	Asteraceae	Sheng	Leaf	Boiled	Chronic cough

of the West Azarbaijan Province, Iran were prepared and compared. There are several reports about the plants which can poses different therapeutic properties and used for several decades in folk medicine. Arasbaran of *Hypericum* were used as digestive, sedative, astringent and strengthening of the respiratory tract, uterine tonic, immune stimulant, anti-depression, anticancer and anti-HIV. The Utica as an astringent and anti-inflammatory effects, is used to fix and discharge of urine. Oregano, in the internal consumption, has the effect of generating power, diuretic, tonic, attenuating of stomach pain, anti-cancer, and relieve migraine pain, and as the external use, it can be beneficial as a rub at the fracture site and numbness in limbs and teeth. It had been reported that Utica, can be used as laxative, anti-cream soda base, stimulant of gastric secretion and peristalsis of the intestines and anticancer[19]. In other regions of Iran, which have traditional religion, the use of medicinal plants is very common even higher than chemical drugs. In Sistan and Baluchistan Province, South-

East of Iran, *Cardaria draba* (L.) Desv, Azmak, was used to soften the breast and relieve anemia[20].

In Shiraz and Kazeroun, both from Fars Province, Southern of Iran, the Armenian marshmallow for colds, bacterial infections, cough, sore throat, angina, bronchitis, and digestive problems are used. Also, Tyghal sugar (*Echinops lalesarensis*) were reported to treat skin and gastrointestinal diseases[21,22].

Traditional use of plants in other province of Iran is also common and this include Kerman[23], Isfahan[24], Ilam[4], and Lorestan[25]. Therapeutic effect of medicinal plants in North, South, East, West and central parts of Iran showed that this country is a vast place to use herbs for treating diseases, however, the use of some plants in different regions of Iran with similar therapeutic effects were observed. Although, each medicinal plant may have its own component to treat special diseases, but, it has been shown that most of these plants possess antioxidant activities[17,26,27]. It should be noted that most of the diseases which are treated with these

medicinal herbs are affected by oxidative stress[28–30]. Oxidative stress has been implicated in respiratory complications and various pathological conditions including aging[31,32], cancer[15,33,34], neurological disorders[35,36], diabetes[37,38], and ischemia/reperfusion injuries[32,38]. These conditions involve many changes, including alterations in the thiol/disulphide redox state, impaired glucose tolerance, and especially, activation of inflammatory processes[39,40]. They are also associated with increased activity of nicotinamide adenine dinucleotide phosphate oxidase or induction of xanthine oxidase with subsequent formation of reactive oxygen species during chronic inflammation and the generation of damage, particularly in respiratory complications[41]. Exposure to agents that increase the formation of reactive oxygen species might accelerate the initiation of pathological changes. Animal models have uncovered a link between an organ's capacity to respond to oxidative stress and diseases development[42,43]. Therefore, plants which have antioxidant activity should be able to counteract these conditions.

In conclusion, traditional medicinal plants can be a source of biological and pharmacological products for the future[44–49]. In general, people believe that the strong positive impact of medicinal plants in the treatment of respiratory disorders. Existed since ancient times, herbs have been used to treat these disorders. Our study revealed the importance of plants in herbal medicine and traditional medicine area.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgements

This research was supported by research grant (No. 453/03) from Deputy for Research and Technology of Lorestan University of Medical Sciences.

### References

- [1] Quiera J. *Basic human histology*. 1st ed. Tehran: Argmand publication; 1993, p. 124–126, 459–485.
- [2] Guyton AC. *Anatomy and physiology*. Philadelphia: Saunders College Publishing; 1985, p. 589–604.
- [3] Asgary S, Sahebkar A, Afshani MR, Keshvari, Haghjooyjavanmard S, Mahmoud Rafieian-Kopaei M. Clinical evaluation of blood pressure lowering, endothelial function improving, hypolipidemic and anti-inflammatory effects of pomegranate juice in hypertensive subjects. *Phytother Res* 2014; **28**(2): 193–199.
- [4] Ghasemi Pirbalouti A, Momeni M, Bahmani M. Ethnobotanical study of medicinal plants used by Kurd tribe in Dehloran and Abadan districts, Ilam Province, Iran. *Afr J Tradit Complement Altern Med* 2013; **10**(2): 368–385.
- [5] Bahmani M, Eftekhari Z. An ethnoveterinary study of medicinal plants in treatment of diseases and syndromes of herd dog in southern regions of Ilam Province, Iran. *Comp Clin Path* 2013; **22**: 403–407.
- [6] Bahmani M, Banihabib E, Rafieian-Kopaei M, Gholami-Ahangaran M. Comparison of disinfection activities of nicotine with copper sulphate in water containing *Limnatis nilotica*. *Kafkas Univ Vet Fak Derg* 2014; doi: 10.9775/kvfd.2014.11223.
- [7] Madihi Y, Merrikhi A, Baradaran A, Rafieian-kopaei M, Shahinfard N, Ansari R, et al. Impact of sumac on postprandial high-fat oxidative stress. *Pak J Med Sci* 2013; **29**(1): 340–345.
- [8] Setorki M, Rafieian-Kopaei M, Merrikhi A, Heidarian E, Shahinfard N, Ansari R, et al. Suppressive impact of *Anethum graveolens* consumption on biochemical risk factors of atherosclerosis in hypercholesterolemic rabbits. *Int J Prev Med* 2013; **4**(8): 889–895.
- [9] Akbari F, Ansari-Samani R, Karimi A, Mortazaei S, Shahinfard N, Rafieian-Kopaei M. Effect of turnip on glucose and lipid profiles of alloxan-induced diabetic rats. *Iran J Endocrinol Metab* 2013; **14**(5): 492–497.
- [10] Nasri H, Rafieian-Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014; **19**(1): 82–83.
- [11] Khosravi-Boroujeni H, Mohammadifard N, Sarrafzadegan N, Sajjadi F, Maghroun M, Khosravi A, et al. Potato consumption and cardiovascular disease risk factors among Iranian population. *Int J Food Sci Nutr* 2012; **63**(8): 913–920.
- [12] Khosravi-Boroujeni H, Sarrafzadegan N, Mohammadifard N, Sajjadi F, Maghroun M, Asgari S, et al. White rice consumption and CVD risk factors among Iranian population. *J Health Popul Nutr* 2013; **31**(2): 252–261.
- [13] Akhlaghi M, Shabanian G, Rafieian-Kopaei M, Parvin N, Saadat M, Akhlaghi M. *Citrus aurantium* blossom and preoperative anxiety. *Rev Bras Anesthesiol* 2011; **61**(6): 702–712.
- [14] Roohafza H, Sarrafzadegan N, Sadeghi M, Rafieian-Kopaei M, Sajjadi F, Khosravi-Boroujeni H. The association between stress levels and food consumption among Iranian population. *Arch Iran Med* 2013; **16**(3): 145–148.
- [15] Azadmehr A, Hajiaghache R, Afshari A, Amirghofran Z, Rafieian-Kopaei M, Yousofi-Darani H, et al. Evaluation of *in vivo* immune response activity and *in vitro* anti-cancer effect by *Scrophularia megalantha*. *J Med Plants Res* 2011; **21**: 2365–2368.
- [16] Shirzad H, Shahrani M, Rafieian-Kopaei M. Comparison of morphine and tramadol effects on phagocytic activity of mice peritoneal phagocytes *in vivo*. *Int Immunopharmacol* 2009; **9**(7–8): 968–970.

- [17] Rafieian–Kopaei M, Baradaran A, Rafieian M. Plants antioxidants: from laboratory to clinic. *J Nephropathol* 2013; **2**(2): 152–153.
- [18] Kafash–Farkhad N, Asadi–Samani M, Rafieian–Kopaei M. A review on phytochemistry and pharmacological effects of *Prangos ferulacea* (L.) Lindl. *Life Sci J* 2013; **10**(8s): 360–367.
- [19] Mirzaei GM, Sewell RDE, Kheiri S, Rafieian–Kopaei M. A clinical trial of the effect of St. John's wort on migraine headaches in patients receiving sodium valproate. *J Med Plants Res* 2012; **6**(9): 1524–1531.
- [20] Iranmanesh M, Najafi S, Yousefi M. Studies on ethnobotany of important medicinal plants in Sistan. *J Herb Drugs* 2010; **1**: 58–65.
- [21] Nasri H, Rafieian–Kopaei M. Medicinal plants and antioxidants: why they are not always beneficial? *Iranian J Public Health* 2014; **43**(2): 255–257.
- [22] Dolatkahi M, Ghorbani–Nahoji M, Mehrafarin A, Amininezhad GR, Dolatkahi A. Ethnobotanical study of medicinal plants in Kazeroon: identification, distribution and traditional uses. *J Med Plants* 2012; **11**(42): 163–178.
- [23] Chaleshtori RS, Rafieian–Kopaei M, Rokni N, Mortezaei S, Sharafati–Chaleshtori A. Antioxidant activity of *Zataria multiflora* hydroalcoholic extract and its antibacterial effect on *Staphylococcus aureus*. *J Mazand Univ Med Sci* 2013; **22**(1): 88–94.
- [24] Mardani–Nejhad S, Vazirpour M. Ethno–botany of medicinal plants by Mobarakeh's people (Isfahan). *J Herb Drugs* 2012; **3**(2): 111–126.
- [25] Ahmadi S, Babakhalo P, Karimifar MA. Medicinal plants of Lorestan. *Yafteh* 2010; **11**(5): 85–100.
- [26] Nasri H, Tavakoli M, Ahmadi A, Baradaran A, Nematbakhsh M, Rafieian–Kopaei M. Ameliorative effect of melatonin against contrast media induced renal tubular cell injury. *Pak J Med Sci* 2014; **30**(2): 261–265.
- [27] Rahzani K, Malekiran AA, Zeraatpishe A, Hosseini N, Seify SM, Abdollahi M. Anti–oxidative stress activity of *Stachys lavandulifolia* aqueous extract in humans. *Cell J* 2013; **14**(4): 314–317.
- [28] Yang FL, Subeq YM, Chiu YH, Lee RP, Lee CJ, Hsu BG. Recombinant human erythropoietin reduces rhabdomyolysis–induced acute renal failure in rats. *Injury* 2012; **43**(3): 367–373.
- [29] Nasri H, Rafieian–Kopaei M. Preventive and curative effect of garlic on nephrotoxic effect of gentamicin in rat. *J Babol Univ Med Sci* 2014; **16**(2): 42–48.
- [30] Yeh PY, Liao FL, Lin SL. Is the renoprotective effect of erythropoietin in chronic kidney disease a myth? *J Formos Med Assoc* 2013; **112**(11): 655–656.
- [31] Rabiei Z, Rafieian–Kopaei M, Mokhtari S, Alibabaei Z, Shahrani M. The effect of pretreatment with different doses of *Lavandula officinalis* ethanolic extract on memory, learning and nociception. *Biomed Aging Pathol* 2013; **4**: 71–76.
- [32] Nasri H, Rafieian–Kopaei M. Oxidative stress and aging prevention. *Int J Prev Med* 2013; **4**(9): 1101–1102.
- [33] Shirzad H, Kiani M, Shirzad M. Impacts of tomato extract on the mice fibrosarcoma cells. *J HerbMed Pharmacol* 2013; **2**(1): 13–16.
- [34] Shirzad H, Tajji F, Rafieian–Kopaei M. Correlation between antioxidant activity of garlic extracts and WEHI–164 fibrosarcoma tumor growth in BALB/c mice. *J Med Food* 2011; **14**(9): 969–974.
- [35] Mirzaei MGR, Azimian M, Moezzi M, Vameghi R, Rafieian–Kopaei M. Effect of lamotrigine on prophylaxis of pediatric classic migraine. *Iran J Child Neurol* 2009; **3**: 35–38.
- [36] Rabiei Z, Rafieian–Kopaei M, Heidarian E, Saghaei E, Mokhtari S. Effects of *Zizyphus jujube* extract on memory and learning impairment induced by bilateral electric lesions of the nucleus basalis of Meynert in rat. *Neurochem Res* 2014; **39**(2): 353–360.
- [37] Mirhoseini M, Baradaran A, Rafieian–Kopaei M. Medicinal plants, diabetes mellitus and urgent needs. *J HerbMed Pharmacol* 2013; **2**(2): 53–54.
- [38] Nasri H, Rafieian–Kopaei M. Protective effects of herbal antioxidants on diabetic kidney disease. *J Res Med Sci* 2014; **19**(1): 82–83.
- [39] Asgary S, Keshvari M, Sahebkar A, Hashemi M, Rafieian–Kopaei M. Clinical investigation of the acute effects of pomegranate juice on blood pressure and endothelial function in hypertensive individuals. *ARYA Atheroscler* 2013; **9**(6): 326–331.
- [40] Ahmadi–Renani S, Fasihi–Ramandi M, Ahmadi K. *Ganoderma lucidum*: a promising anti–inflammatory medicinal plant. *J HerbMed Pharmacol* 2014; **3**(1): 67–68.
- [41] Rafieian–Kopaei M, Baradaran A, Rafieian M. Oxidative stress and the paradoxical effects of antioxidants. *J Res Med Sci* 2013; **18**(7): 629.
- [42] Sarrafzadegan N, Khosravi–Boroujeni H, Esmailzadeh A, Sadeghi M, Rafieian–Kopaei M, Asgary S. The association between hypertriglyceridemic waist phenotype, menopause, and cardiovascular risk factors. *Arch Iran Med* 2013; **16**(3): 161–166.
- [43] Setorki M, Nazari B, Asgary S, Azadbakht L, Rafieian–Kopaei M. Anti atherosclerotic effects of verjuice on hypocholesterolemic rabbits. *Afr J Pharm Pharmacol* 2011; **5**(8): 1038–1045.
- [44] Bahmani M, Abbasi J, Mohsenzadegan A, Sadeghian S, Gholami–Ahangaran M. *Allium sativum* L.: the anti–immature leech (*Limnatis nilotica*) activity compared to Niclosomide. *Comp Clin Pathol* 2013; **22**: 165–168.
- [45] Amirmohammadi M, Khajoenia S, Bahmani M, Rafieian–Kopaei M, Eftekhari Z, Qorbani M. *In vivo* evaluation of antiparasitic effects of *Artemisia abrotanum* and *Salvia officinalis* extracts on *Syphacia obvelata*, *Aspiculuris tetrapetra* and *Hymenolepis nana* parasites. *Asian Pac J Trop Dis* 2014; **4**(Suppl 1): S250–S254.
- [46] Bahmani M, Farkhondeh T, Sadighara P. The anti–parasitic effects of *Nicotina tabacum* on leeches. *Comp Clin Pathol* 2012; **21**(3): 357–359.
- [47] Bahmani M, Rafieian–Kopaei M. Medicinal plants and secondary metabolites for leech control. *Asian Pac J Trop Dis* 2014; **4**(4): 315–316.
- [48] Ghasemi–Pirbalouti A, Bahmani M, Avijgan M. Anti–*Candida* activity of some of the Iranian medicinal plants. *Electron J Biol* 2009; **5**(4): 85–88.
- [49] Bahmani M, Saki K, Rafieian–Kopaei M, Karamati SA, Eftekhari Z, Jelodari M. The most common herbal medicines affecting Sarcomastigophora branches: a review study. *Asian Pac J Trop Biomed* 2014; **4**(12): 930–937.